

AD-A075 958

OKLAHOMA UNIV NORMAN DEPT OF CHEMISTRY

VARIABLE TEMPERATURE 119M SN MOESSBAUER STUDY OF ORGANOTIN - SU--ETC(U)

OCT 79 K C MOLLOY , J J ZUCKERMAN

N00014-77-C-0432

F/G 11/9

UNCLASSIFIED

TR-13

NL

| OF |

AD
A075958




END

DATE

FILMED

11-79

DDC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

UNCLASSIFIED

LEVEL

B.S. 12

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE

READ INSTRUCTIONS
BEFORE COMPLETING FORM

1. REPORT NUMBER 13	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Variable Temperature ^{119m} Sn Mössbauer Study of Organotin - Substituted Styrene Monomers and Polymers.		5. TYPE OF REPORT & PERIOD COVERED
7. AUTHOR(s) K. C. Molloy, J. J. Zuckerman, H. Schumann, G. Rodewald		8. CONTRACT OR GRANT NUMBER(s) N00014-77-C-0432
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Oklahoma Department of Chemistry Norman, Oklahoma 73019		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 053-636
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy Arlington, Virginia 22217		12. REPORT DATE 1 Oct 79
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 12 8 14 TR-13		13. NUMBER OF PAGES 6
15. SECURITY CLASS. (of this report) Unclassified		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for Public Release, Distribution Unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) Prepared for Publication in Inorganic Chemistry		
18. SUPPLEMENTARY NOTES 9 Technical rept's		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Organotin-Substituted Styrenes, Styrene Polymers, Mössbauer Spectroscopy, Tin- ^{119m} Mössbauer Spectroscopy, Variable Temperature Mössbauer Spectroscopy, Lattice Dynamics, Polymerization.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Variable temperature ^{119m} Sn Mössbauer spectroscopy can detect the enhanced binding of the tin atom in organotin(IV)-substituted styrenes on polymerization.		

DDC
OCT 31 1979
NEGATIVE
D

79 10 31 030

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-LF-014-6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

403 980

alt

A 075958

DDC FILE COPY

OFFICE OF NAVAL RESEARCH

Contract N00014-77-C-0432

Task No. NR 053-636

TECHNICAL REPORT NO. 13

Variable Temperature ^{119}Sn Mössbauer Study of Organotin -
Substituted Styrene Monomers and Polymers.

by

K. C. Molloy and J. J. Zuckerman
H. Schumann and G. Rodewald

Prepared for Publication

in


Inorganic Chemistry

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DDC TAB	<input checked="checked" type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	<input type="checkbox"/>
By _____	
Distribution/	
Availability Codes	
Dist.	Avail and/or special
A	

University of Oklahoma
Department of Chemistry
Norman, Oklahoma 73019

Reproduction in whole or in part is permitted for
any purpose of the United States Government

*This document has been approved for public release
and sale; its distribution is unlimited



Variable Temperature $^{119}\text{m}\text{Sn}$ Mössbauer Study of Organotin -
Substituted Styrene Monomers and Polymers.

K. C. Molloy and J. J. Zuckerman
Department of Chemistry
University of Oklahoma
Norman, OK, 73019, USA

and

H. Schumann and G. Rodewald
Institute for Inorganic and Analytical Chemistry
Technical University of Berlin
1 Berlin 12 (Charlottenburg) GERMANY

SUMMARY

Variable temperature $^{119}\text{m}\text{Sn}$ Mössbauer spectroscopy can detect the enhanced binding of the tin atom in organotin(IV)-substituted styrenes on polymerization.

Tin-containing organometallic polymers (OMPs) are being actively investigated as potential marine antifouling materials. The OMPs are unique in containing the organotin pesticide chemically bound to a polymer backbone, resulting in controlled release into the marine environment.¹ The binding at the tin site is crucial to the rate at which the release will take place. We report here the first application of a method based upon Mössbauer spectroscopy for distinguishing the state of binding of the tin atoms in organotin-substituted monomers and polymers.

Di-para-totyldi-para-styryltin(IV) (I) (mp 83-84°C) and tetra-para-styryltin(IV) (II) (mp 134.5-135.5°C) are solids formed by treatment of the corresponding tin chloride with the para-styryl Grignard reagent. The monomers are polymerized to (Ia) and (IIa) by gentle heating to infusible solids.² The tin-119m Mössbauer Isomer Shift (IS) and Quadruple Splitting (QS) values for the monomers (I: IS=1.23±0.02; QS=0mm/s and II: IS=1.17±0.02; QS=0mm/s) do not change outside of experimental error on polymerization (Ia: IS=1.21±0.02; QS=0mm/s and IIa: IS=1.20±0.02; QS=0mm/s).³ These values are typical of simple tetraorganotin compounds,⁴ and the failure to detect any change in their magnitudes on polymerization has been noted before for trimethyl-⁵ and triphenyl-para-styryltin(IV).⁶

It has been recently found that logarithmic plots of the Mössbauer resonance areas vs. temperature are linear over an accessible temperature range as expected according to the Debye model of solids in the high temperature limit. For a thin absorber the recoil-free fraction, $f(T)$, is directly related to this area, $A(T)$, and may be used to investigate the lattice dynamics at the tin site. It is found that the more tightly bound the tin atoms, the slower will be the decrease in f , and hence $A(T)$, as the temperature is raised. Compounds of known structure, consisting of non-interacting monomeric molecules exhibit slopes $[a = \frac{d \ln (A_T/A_{77})}{dT}]$ of ca. $-1.8 \times 10^{-2} K^{-1}$. Solids in which one-, two- or three- dimensional association is present exhibit slopes of ca. $-0.9 \times 10^{-2} K^{-1}$ or less.⁷

Figure 1 displays the logarithmic plot of the area under the Mössbauer resonances, normalized to 77K for ease of comparison of data, vs. temperature

in the range 77 to 140K for monomers I and II and their polymers Ia and IIa.

The spectral intensities in the case of the monomers decays rapidly [I: $a = -2.13 \times 10^{-2} \text{ K}^{-1}$ (regression analysis, $r = -0.990$, intercept 1.64, 4 points); II: $a = -2.08 \times 10^{-2} \text{ K}^{-1}$ ($r = -0.998$, 1.60, 3)], and it becomes impossible, even over extremely long counting times, to obtain usable data above ca. 100K. By contrast the spectral intensities for the materials after polymerization show slower decays [Ia: $a = -1.79 \times 10^{-2} \text{ K}^{-1}$ ($r = -0.993$, 1.36, 8); IIa: $a = -1.95 \times 10^{-2} \text{ K}^{-1}$ ($r = -0.999$, 1.51, 6)], and spectra can be obtained up to ca. 140K.

The effect of polymerization in locking the tin atom in its position must necessarily be very small, yet the effect is detectable by Mössbauer techniques in two ways. Firstly, the temperature dependences of the recoil-free fractions are discernably greater for the monomers than for their polymers. Secondly, the Debye model by which the slopes in Figure 1 are linear, itself begins to break down, and for the monomers the resonance areas above ca. 100K are very much smaller than expected by extrapolation of the linear plots in the range 77 to 100K.

While the tin atoms are clearly more free to move in the monomers, the added constraint on polymerization is not as marked as when tin atoms are bound rigidly by bridging, directly-attached electronegative substituents.⁷ In the case of the tetraorganotins studied here, the tin atoms continue to be held by bonds to four carbons and, moreover, the polymerized lattice itself may itself be quite porous owing to the steric bulk of the atoms attached to tin.

Acknowledgement.

Our work is supported by the Deutsche Forschungsgemeinschaft, Bonn-Bad Godesberg and the Fonds der Chemischen Industrie, Frankfurt (to H.S.) and by the Office of Naval Research (to J.J.Z.). We thank especially the Scientific Affairs Division of NATO for a grant which makes this collaboration possible.

References and Notes

- (1) Yeager, W.L.; Castelli, V.J.; Abstr. 173rd American Chemical Society National Meeting, New Orleans, LA, March, 1977; Subramanian, R.V.; Garg, B.K. and Corredor, J.; in Organometallic Polymers, ed. by Carraher, C.E.; Academic Press, New York, 1978, p. 181; Subramanian, R.V.; Garg, B.K.; Polym.-Plast. Technol. Eng. 1978, 11, 81.
- (2) Rodewald, G.; Doctoral Dissertation, Technical University of Berlin, 1978.
- (3) Schumann, H.; Rodewald, G.; Rodewald, V.; Lefferts, J.L. and Zuckerman, J.J.; submitted for publication.
- (4) Zuckerman, J.J.; Adv. Organometal. Chem., 1970, 9, 21; Ruddick, J.N.R.: Revs. Silicon, Germanium, Tin Lead Comps., 1976, 2, 115.
- (5) Aleksandrov, A. Yu.; Okhlobystin, O. Yu.; Polak, L.S.; Shpinel, V.S.; Dokl. Akad. Nauk SSSR 1964, 157, 934.
- (6) Khrapov, V.V.; Goldanskii, V.I.; Prokofiev, A.K.; Kostyanoviskii, R.G.; J. Gen. Chem. USSR 1967 37, 1.
- (7) Harrison, P.G.; in Organotin Compounds: New Chemistry and Applications, ed. by Zuckerman, J.J.; Advances in Chemistry Series No. 157, American Chemical Society, Washington, D.C., 1976, p. 258; Harrison, P.G.; Phillips, R.C.; Thornton, E.W.; J. Chem. Soc., Chem. Commun. 1977, 603; Lefferts, J.L.; Molloy, K.C.; Zuckerman, J.J.; Haiduc, I.; Guta, C.; Ruse, D. submitted for publication.

Figure Caption

Figure 1. Logarithmic plot of the area under the Mössbauer resonances $[A(T)]$ normalized to 77K vs. temperaturer in the range 77 to 140K for Di-para-tolyl-di-para-styryltin(IV) and tetra-para-styryltin(IV) and their polymers.

$\text{Log}_e[A(T)/A(77)]$

